



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2020

The geography of speciation in dasyurid marsupials

García-Navas, Vicente ; Kear, Benjamin P ; Westerman, Michael

Abstract: Aim To determine the effects of competition and divergence time on morphological dissimilarity and geographical range overlap between dasyurid species at both regional and local scales. Our hypothesis is that speciation in this group has been largely allopatric at regional scale, but involved morphological divergence at local scale through sympatric character displacement. Location Australia, New Guinea and surrounding islands. Taxon Dasyurid (Dasyuridae) marsupials, 67 species. Methods Geographical range overlap was quantified using polygons representing the outer limits of species distributions. Local-scale range overlap was quantified as the degree of co-occurrence of two taxa across a set of ecological survey plots representing 83 sampled communities. Phylogenies were generated using a novel DNA dataset, with divergence times estimated via total-evidence dating incorporating fossils. Morphological divergence was determined using body mass and lower molar row length as proxy traits for reconstructing niche exploitation. Results Sister species pairs were found to be sympatric in 52% (11/21) of cases. Range overlap tended to increase with node age, which supports the hypothesis that mammalian speciation is routinely allopatric. We detected no evidence of character displacement with increasing range overlap between sister species pairs. However, a negative relationship was observed between morphological divergence in body mass and range overlap across all sampled taxa, suggesting that selection in sympatry is convergent, while divergent selection occurs in allopatry. Local-scale co-occurrences revealed no trace of species aversion, indicating that competition has not impacted on the spatial distribution of dasyurids. Main conclusions Despite moderate levels of sympatry through time, our results evince low rates of spatial co-occurrence between dasyurid species. Although this may be indicative competitive exclusion, the lack of character displacement suggests that biotic interactions have likely not acted as a dominant driver of phenotypic evolution in this radiation. We alternatively posit that abiotic factors including aridity and geographical connectivity have more feasibly propagated character convergence, and led to both niche conservatism and speciation in this ubiquitous australidelphian clade.

DOI: <https://doi.org/10.1111/jbi.13852>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-198876>

Journal Article

Accepted Version

Originally published at:

García-Navas, Vicente; Kear, Benjamin P; Westerman, Michael (2020). The geography of speciation in dasyurid marsupials. *Journal of Biogeography*, 47(9):2042-2053.

DOI: <https://doi.org/10.1111/jbi.13852>

The geography of speciation in dasyurid marsupials

Vicente García-Navas^{1*}, Benjamin P. Kear² & Michael Westerman³

¹ *Department of Evolutionary Biology and Environmental Studies, University of Zurich, Zurich, Switzerland*

² *Museum of Evolution, Uppsala University, Uppsala, Sweden*

³ *Department of Environment, Ecology and Evolution, La Trobe University, Melbourne, Victoria, Australia*

**Correspondence:* Vicente García-Navas - Department of Evolutionary Biology and Environmental Studies, University of Zurich, Winterthurerstrasse 190, 8057, Zurich, Switzerland.

Tel. +41 (0) 44 63 54914. E-mail: vicente.garcianavas@gmail.com

ORCID:

Vicente García-Navas 0000-0002-9362-2663

Benjamin P. Kear 0000-0002-3128-3141

Running head: Marsupial range overlap and trait divergences

Word count (excl. Suppl. Mat): 5,187

Number of references: 66

Number of figures: 3

Number of tables: 0

37 *Abstract*

38 *Aim:* To determine the effects of competition and divergence time on morphological
39 dissimilarity and geographic range overlap between dasyurid species at both regional and local
40 scales. Our hypothesis is that speciation in this group has been largely allopatric at regional
41 scale, but involved morphological divergence at local scale through sympatric character
42 displacement. *Taxon:* Dasyurid (Dasyuridae) marsupials, 67 species. *Location:* Australia, New
43 Guinea and surrounding islands. *Methods:* Geographic range overlap was quantified using
44 polygons representing the outer limits of species distributions. Local-scale range overlap was
45 quantified as the degree of co-occurrence of two taxa across a set of ecological survey plots
46 representing 83 sampled communities. Phylogenies were generated using a novel DNA dataset,
47 with divergence times estimated via total-evidence dating incorporating fossils. Morphological
48 divergence was determined using body mass and lower molar row length as proxy traits for
49 reconstructing niche exploitation. *Results:* Sister species pairs were found to be sympatric in
50 52% (11/21) of cases. Range overlap tended to increase with node age, which supports the
51 hypothesis that mammalian speciation is routinely allopatric. We detected no evidence of
52 character displacement with increasing range overlap between sister species pairs. However, a
53 negative relationship was observed between morphological divergence in body mass and range
54 overlap across all sampled taxa, suggesting that selection in sympatry is convergent, while
55 divergent selection occurs in allopatry. Local-scale co-occurrences revealed no trace of species
56 aversion, indicating that competition has not impacted on the spatial distribution of dasyurids.
57 *Main conclusions:* Despite moderate levels of sympatry through time, our results evince low
58 rates of spatial co-occurrence between dasyurid species. Although this may be indicative
59 competitive exclusion, the lack of character displacement suggests that biotic interactions have
60 likely not acted as a dominant driver of phenotypic evolution in this radiation. We alternatively
61 posit that abiotic factors including aridity and geographic connectivity have more feasibly
62 propagated character convergence, and led to both niche conservatism and speciation in this
63 ubiquitous australidelphian clade.

65 *Keywords:* age-range correlation; allopatric speciation; Australia; community phylogenetics;

66 Dasyuridae; sister species

67

68

1. INTRODUCTION

Explaining patterns of species co-occurrence in relation to their ecological and morphological characteristics constitutes a core goal in many ecological investigations. Indeed, over the last 15 years, the incorporation of phylogenetic data into classic ecological studies on species coexistence has contributed to illuminate the processes underlying biotic assembly at different levels (from a local to a regional perspective) (reviewed in Warren et al., 2014). In this sense, two distinct approaches have been commonly employed, each one of them focused on a different spatial scale. On one hand, site-based approaches quantify spatial variation as the relatedness between co-occurring taxa and include measures of community phylogenetic structure and phylogenetic diversity (Webb et al., 2002; Vamosi et al., 2009). These methods - grouped into the field of community phylogenetics- are normally used to determine the role of ecological mechanisms (e.g., competition or environmental filtering) in structuring communities (Cavender-Bares et al., 2009; Cantalapiedra et al., 2014). On the other hand, regional scale analyses compare spatial distribution at deeper clade levels, and can thus reveal broader drivers of geographical speciation by quantifying the extent of range overlap between sister lineages (Anacker & Strauss, 2014; Hodgel & Bellwood, 2016; Gaboriau et al., 2018). Although local and regional scale analyses should ideally be applied in tandem to mitigate the influence of historical biogeography on currently extant species distributions (Mouquet et al., 2012; Wiens, 2012), only a few studies have evaluated such interlinked ecological and evolutionary perspectives simultaneously (Borregaard et al., 2014; Weeks et al., 2016). This paucity of studies addressing simultaneously the roles of history and ecology in biotic assembly is noteworthy as it is fairly clear that speciation history can impinge on community patterns at local scales (Cardillo, 2012), and likewise, there is evidence for the imprint of competition on broad-scale assemblage structure (Cardillo, 2011). Hence, although species with minimal regional overlap will usually exhibit limited co-occurrence at community level, the converse does not have to be necessarily true as fine-scale mechanisms (e.g., competitive exclusion) could prevent the coexistence of two species even when they show overlapping distributions. Consequently, under certain circumstances a high degree of range overlap may not translate into

real coexistence at a local scale. That is, in many cases sympatric species (in terms of broad-scale spatial overlap) are not syntopic and do not interact ecologically (Cardillo & Warren, 2016).

Here, we assess the patterns of local and regional scale species co-occurrence within Dasyuridae, a radiation of small-medium-sized carnivorous marsupials that incorporate some 75 named species (van Dyck & Strahan, 2008). Dasyurids are distributed across virtually every terrestrial environment throughout Australasia, ranging from the Central Australian arid zone to tropical rainforests in northeastern Australia and New Guinea (Dickman, 2003). The Dasyuridae constitute a geologically recent radiation, which originated around the Palaeogene-Neogene boundary (approximately 23 Ma), but diverged into its major crown groupings across the Mid-Miocene Climatic Optimum (beginning around 18 Ma); this coincides with the tectonic connection of Australia with New Guinea and the onset of cooler, dryer climatic conditions that propagated more open habitats and intracontinental aridity (Krajewski et al., 2000; Westerman et al., 2016; Kealy and Beck, 2017). These landmark events undoubtedly created novel ecological niches, probably as result of the decline of competitors (e.g., thylacinids or ‘marsupial wolves’, and peramelemorphians, also known as bandicoots), and/or the emergence of new habitats that promoted dispersal and vicariance processes (Morton et al., 1989). As result of the vicariance speciation events, many extant dasyurid species today occur in only a narrow range of habitat types, or over restricted geographical distributions and show allopatric distributions. For example, the widespread xeric-adapted Dunnart, *Sminthopsis crassicaudata*, is usually associated with cracking clay soils, whereas its close relative *Sminthopsis douglasi* is limited to alluvial plain Mitchell grass pastures in central-northern Queensland (Woolley, 1992). Likewise, spinifex (*Triodia*) grasses have been integral to the radiation of many arid-zone dasyurid lineages (Anderson et al., 2019), including the tiny rodent-like Ningauis (*Ningauis*), which primarily inhabit spinifex hummocks (Dickman, 2003).

Although they are widely distributed the continent and can be found in a wide range of habitats, dasyurids exhibit a marked lack of morphological specialization compared to other australidelphian marsupials (van Dyck & Strahan, 2008). Dasyurid marsupials show a

constrained basic body-plan and, apparently, morphological diversification has mostly involved changes in body size (from mouse-like to dog-sized species), seemingly related with dietary preferences: most dasyurids are insectivorous, except for the Mulgara (*Dasycercus cristicauda*), Kowari (*Dasyuroides byrnei*), largest (>500 g) quolls (*Dasyurus*), and the Tasmanian devil, all of which feed upon vertebrates (Fisher & Dickman, 1993). The lack of feeding niche segregation amongst dasyurids suggests that the partitioning of food resource appears to occur on prey size alone, which would accentuate the associated effect on body size differences (Fox, 1982). In this regard, Jones (1997) found evidence for competitive character displacement in body size (measured as skull length) within the guild of dasyurid carnivores inhabiting Tasmania (see also Jones & Barmuta, 1998). Subtler morphological and behavioural differences appear to be common amongst sympatric species of the smaller-bodied insectivorous lineages (Dickman, 1984). For example, the long-tailed (*Planigale ingrami*), and narrow-nosed (*Planigale tenuirostris*) planigales, which maintain overlapping distributional ranges, can be distinguished by relative flattening of the cranium (Archer, 1976). Intraspecific competition has also been evoked for explaining why the agile antechinus (*Antechinus agilis*), a scansorial species, spend more time foraging on the ground once the sympatric and fossorial Swainson's antechinus (*A. swainsonii*) is removed from its range (Dickman, 1986a,b). A similar phenomenon has been proposed (Dickman, 1988) to explain the delayed breeding schedule of *A. agilis* in comparison to its co-occurring relative, the brown or Stuart's antechinus (*A. stuartii*).

Noticeably, the strength of competition appears to be greater in mesic environments that sustain high population densities (forests, heathlands) but where the number of co-occurring species is low (2-3 spp.) in comparison with xeric habitats, where the unpredictability of food resources maintains population densities at low levels, thereby decreasing the probability of interspecific encounters (Dickman, 2003). Despite the extreme climatic conditions (variable rainfall, severe drought) to which organisms present in arid and semi-arid environments must face, up to 9 dasyurid species can co-exist in this region. The unexpected high dasyurid diversity present in harsh environments (deserts, hummock grasslands) characteristics of the Australian arid zone has been associated to the existence of increased cladogenesis in this region

in several taxonomic groups including the Dasyuridae (e.g., Rabosky et al., 2007; Catullo & Keogh, 2014; García-Navas et al., 2018). At a local scale, lack of pressure from competition may also have contributed to maintain high species richness in this region (Crowther & Blacket, 2003). Alternatively, such patterns may arise as result of the existence of substantial niche partitioning based on body size (i.e., small body size overlap) in agreement with the limiting similarity principle (MacArthur & Levins, 1967; Grant & Grant, 2006).

In this study, we examine the relationship between species co-occurrence patterns and morphological similarity in dasyurids at both regional and local (community) scales. Our hypothesis is that speciation in this group has been largely allopatric at regional scale, as previously shown in other taxonomic groups (Fitzpatrick & Turelli, 2006), and that in those cases where sympatry exists, it involves morphological divergence at local scale through character displacement (e.g., Grossenbacher & Whittall, 2011). Although interspecific competition should favor diversification of functional traits, competition for resources may also lead to adaptive convergence whether the benefits to an individual of excluding heterospecifics from their territory are similar to the benefits of excluding conspecifics (Cody, 1969; Drury et al., 2018). This is especially pertinent for harsh xeric environments where density-dependent factors are important, and selection may favor trait convergence (e.g., body mass) to establish dominance hierarchies and mediate territorial interactions (e.g., Fox, 1982). Thus, we controlled for similarity in habitat when testing whether interspecific competition leads to morphological convergence. Next, we used community data to address the relationship between local scale species co-occurrence and morphological divergence and test whether those species that co-exist in the same assemblages (positive associations) are more morphologically divergent than those that do not coexist (negative associations) or do so not preferentially (random associations). Lastly, we used a novel approach, node-based analysis (Borregaard et al., 2014), to quantify the distributional divergence between daughter lineages descending from the same node and thereby obtaining a better understanding of the phylogenetic structure of species assemblages. Since the four tribes conforming this family share a complex biogeographic

history of dispersals across Australia and Papua New Guinea (García-Navas et al., 2018), we expect a low number of nodes to exhibit strong distributional change.

2. MATERIAL AND METHODS

2.1 Phylogeny and sister species pair selection

We first derived a novel phylogeny of Dasyuridae using a modified version of the total-group Dasyuromorphia DNA sequence dataset compiled by Westerman *et al.* (2016). This 17,968 bp matrix included an expanded ingroup sample of 80 extant dasyurid taxa (see Fig. 1), together with nDNA sequence data for the omega globin, runt-related transcription factor 2 (*RUNX2*), and *col2A1* enhancer genes (see Supplementary Data). Corresponding information was also transcribed from the recently published outgroup thylacine (*Thylacinus cynocephalus*) genome (Feigin et al., 2018). We then ran a subset analysis that pruned the ingroup dasyurid taxon sample (primarily via deletion of closely related terminals within the *Antechinus* species complex) to match the 67 extant species represented by our ecomorphological data. A phylogenetic divergence timescale was estimated using node + tip total evidence dating following Kealy & Beck (2017). We augmented their 115 craniodental/postcranial skeletal character dataset with five new fossil taxa to test alternative node calibrations (total taxon sample = 100), but otherwise retained the original parameters and settings. Further details about our phylogenetic and tree dating analyses are provided in the Supplementary Material.

Twenty-one sister species pairs (= most closely related sister taxa) were identified from our time-calibrated total evidence tree based on >0.95 posterior probability (PP) support at their most recently divergent node (see Supplementary Material). Because the majority of taxa not included in our analysis belonged to the same *Antechinus* species complex, we deemed their omission to have minimal impact on our overall hypothesis of dasyurid interspecific relationships. Pairwise divergence times were then calculated using the ‘chronoMPL’ function in the *ape* library (Paradis et al., 2004).

2.2. Character selection and divergence estimation

We gathered data on overall body size (expressed as both body mass and maximum snout-tail length) and length of the lower molar row (m_1 - m_4) for a sample set of 67 dasyurid species (accounting for 89% of the total clade species count) based on information available in the published literature (Table S1; Supplementary Data). Overall body size, which ranges from 52 mm and 11.6 g in *Ningaui yvonneae* to 611 mm and 8,000 g in *Sarcophilus harrisii*, constitutes a key attribute for any organism determining ecological interactions (e.g., who eats whom and at what rate) and confers fitness advantages in form of size-dependent competition for resources, increased fecundity and greater diet breadth (e.g., Kraaijeveld-Smit et al. 2003; Fisher & Cockburn, 2006). As an example, Dickman (2014) reported a positive relationship between body mass and percentage intake of vertebrates in the diet of 25 dasyurid species. Dental morphology is also generally indicative of diet (e.g., Pineda-Munoz et al., 2017; Martin et al., 2016), and while dasyurids lack specialized carnassials, their four pairs of molars are massive and structurally comparable between species (Archer, 1976; 1978). Because overall body and tooth size can be correlated in marsupials (Gordon, 2003), the length of the lower molars was corrected for body size using a least squares regression analysis. The resulting residuals were used as input in further analyses. As body size and body mass are highly correlated ($r = 0.85$, $p < 0.001$) and we obtained similar results, for the sake of brevity we only report the results of our body mass analyses here.

We used sister species pair comparisons to evaluate the relationship between morphological disparity in overall body size and molar row length against patterns of regional co-occurrence. Character divergences were quantified following the approaches of Davies et al. (2007) and Anacker & Strauss (2014), which contrast the logarithm of trait values between sister-pairs in the form of: $\log(X_A) - \log(X_B)$ where X was the trait value and $X_A > X_B$.

2.3 Range overlap estimation

To quantify range overlap between species pairs, we converted species' distribution maps retrieved from the IUCN website (IUCN, 2018) to vector polygons. These were collated with

official biodiversity records from the Atlas of Living Australia (<http://www.ala.org.au>), and rasterized to a 100×100 km grid using ArcGIS to obtain a presence-absence matrix. This dataset was used to compute the level of range overlap between species pairs using the R package *letsR* (Vilela & Villalobos, 2015). Range overlap represents the proportion of co-occurrence affecting each species within the combined range of each pair of species (Chesser & Zink, 1994; Martin et al., 2010); this is expressed as a value between 0 (no overlap) and 1 (individual species range completely overlapped by the larger range). We accordingly classified our 21 sister species pairs as being either allopatric (no overlap), or sympatric (overlap), and examined the variation in range overlap relative to divergence time and degree of morphological dissimilarity (evidenced by contrasting phenotypic traits) using univariate regression models. To test the robustness of our results against models of trait evolution (Brownian motion: Felsenstein, 1985), we repeated these regressions with correction for divergence time by dividing the observed overlap with the square root of the sum of the branch lengths connecting the species pairs (*sensu* Davies et al. 2007). In addition, we assessed morphological dissimilarity between allopatric and sympatric pairs of sister species by applying *t*-tests.

2.4 Geography of speciation

We examined how range overlap varies with divergence time using an ‘age-range correlation’ (ARC) test to yield information about the geography of speciation (see Fitzpatrick & Turelli, 2006). Intercept values ≥ 0.5 and a negative slope were treated as indicative of predominantly sympatric speciation, whereas intercepts ≤ 0.5 and positive slopes are suggestive of allopatric speciation (Fitzpatrick & Turelli, 2006).

2.5 Testing for sympatric convergence

We created a dissimilarity index by calculating the square-root transformed Euclidean phenotypic distance based on differences in terms of body mass and molar row length between all pairs of species. This was then correlated with our pairwise matrix of range overlap using a Mantel test to establish whether sympatry is negatively associated with dissimilarity, as would

be expected if trait convergence arises as consequence of interspecific interactions. Because convergence in sympatry can also result from selection favoring particular phenotypes from comparable environments, we accommodated for habitat by using generalized categorizations: 1. forests; 2. woodlands, heathlands and mallee; and 3. grasslands and sand-plains. If, after controlling for compatibility in environmental conditions, sympatry is negatively correlated with the morphological dissimilarity index, then this would support the inferred existence of trait convergence as result of interspecific interactions. We used partial Mantel tests to examine the relationship between spatial overlap and morphological divergence controlling for phylogenetic distance. Patristic distances were computed using *adephylo* (Jombart et al., 2010). Mantel tests were implemented in R using the *vegan* package (Oksanen, 2010).

2.6 Local scale species co-occurrence

To assess whether our observed regional scale patterns are thus informative of processes operating at local scales (e.g., ecological interactions), we quantified local-scale co-occurrences between Australian species. We compiled occurrence data for a total of 83 local communities from scientific articles, book chapters, and supplements of the Western Australian Museum (see Supplementary References for full literature list). We then tested for statistically significant pairwise patterns of species co-occurrence by applying the approach proposed by Veech (2013) to our community data-based presence-absence matrix. Thereby, we determined the probability that any pair of species would co-occur at a frequency greater or lesser than the observed frequency if the two given species are distributed randomly (independently) of one another. We only considered those species pairs that co-occurred across a minimum number of plot surveys, and yielded frequencies >1 . We then tested for differences in the level of morphological similarity between those species that positively co-occurred, and those that showed a negative or random association.

In addition, we quantified the degree of co-occurrence among all pairs of species across our local dataset (i.e., Australian taxa that were recorded in at least one survey plot) *via* their *c*-scores (= checkerboard score: Stone & Roberts, 1990). High *c*-scores indicate reduced instances

of co-occurrence, and suggest the influence of mechanisms such as competitive exclusion. Potential relationships between local and regional scale co-occurrences (measured as degree of range overlap and *c*-scores, respectively), phylogenetic distance, and level of morphological dissimilarity in sympatric species were further assessed using Mantel tests with non-overlapping species pairs excluded. These analyses were implemented using the R packages *cooccur* (Griffith et al., 2016) and *picante* (Kembel et al., 2010).

2.7 Node-based analysis

We used the node-based method of Borregaard et al. (2014) to combine ecological and evolutionary patterns into a single framework; this samples each node in a given phylogeny and compares the distribution of descendant clades in our phylogeny against a null model. The resulting ‘specific overrepresentation score’ (SOS) provides a measure of clade overrepresentation that integrates both clade-based and site-based approaches. SOS values calculated for a certain node can be summarized across occupied sites to yield the ‘geographic node divergence’ (GND), which quantifies the distributional divergence between the two daughter lineages descended from a given node. GND thus identifies which nodes are responsible for observed patterns of phylogenetic structure and regional scale species co-occurrence (see Supplementary Material for details).

3. RESULTS

3.1 Phylogenetic and tree dating inferences

Our analysis of the expanded DNA sequence phylogenetic dataset yielded an almost identical tree topology (Fig. S1) to that produced by Westerman et al. (2016). All nodes comprising the extant ingroup dasyurid clades retrieved virtually unanimous support (PP = 0.98–1), except for placement of the dunnart species *Sminthopsis granulipes* (PP = 0.77), and *Sminthopsis youngsoni* + *Sminthopsis hirtipes* (PP = 0.61) within Sminthopsini, and the planigales *Planigale*

maculata + *Planigale novaeguineae* + *Planigale ingrami* + *Planigale gilesi* (PP = 0.57), and *P. ingrami* + *P. gilesi* (PP = 0.7) within Planigalini (compare with Westerman et al., 2017).

Our total evidence node + tip dated tree topology (Fig. 1) generated dasyurid crown clade divergence estimates (Table S2) that are generally younger than those advocated by other studies (Westerman et al., 2016; Kealy & Beck, 2017). Most significantly, the initial radiation of extant dasyurid lineages was reconstructed as coinciding with the Mid-Miocene Climatic Optimum (around 15 Ma), a result that post-dates other estimates by up to 5 Ma (see Kealy & Beck, 2017). Cladogenic events propagating the extant dasyurid species-level lineages thus accompany the onset of aridity during the late Miocene and Pliocene (commencing from about 10 Ma). However, we emphasize that divergence times within Dasyuromorphia are sensitive to a priori dataset selection and preferred parameters, with DNA-only analyses consistently yielding substantially older node ages (see discussion in Kealy & Beck, 2017); this concurs with the contrasting time-trees generated for other australidelphian marsupial clades (e.g., Kear et al., 2016; Travouillon & Philips, 2018).

3.2 Sister species analyses

We detected no statistically significant differences in average node age and divergence in terms of body mass or molar row length between sympatric and allopatric sister species pairs within Dasyuridae (*t*-tests, all *p*-values > 0.1; see Table S3). The degree of range overlap between these same species pairs was also not correlated with the level of morphological divergence regardless of which trait was analyzed (body mass: $F = 0.16$, $p = 0.70$; molar row length: $F = 0.01$, $p = 0.92$). We obtained similar results when repeating the regression models controlling for divergence time (both *p*-values > 0.5). There was likewise no significant linear relationship between divergence time and either body mass ($F = 0.38$, $p = 0.25$) or molar row length ($F = 0.11$, $p = 0.74$), even when analyzing sympatric and allopatric pairs separately (all *p*-values > 0.05).

3.3 Age-range correlation test

Range overlap tends to increase with time since divergence (node age), although this relationship was not statistically significant (slope = 0.07, intercept = 12.47, $F = 1.39$, $p = 0.25$). This result was also consistent (slope = 0.06, intercept = 10.51, $F = 1.25$, $p = 0.28$) after excluding sister species pairs with PP values <1 (Fig. 1).

3.4 Range overlap and morphological dissimilarity

We found no significant relationship between the level of sympatry (range overlap; average value = 11.8%) and morphological distance between species pairs (Mantel test, $p = 0.11$). However, pairwise dissimilarity in body mass was negatively correlated with the degree of range overlap (Mantel test, $r = 0.06$, $p = 0.015$; Fig. 2) when examined on its own; this suggests that species more similar in terms of body mass overlap more in their geographical range. When grouped by habitat type, a relationship close to being significant was also identified for species living in grasslands and sandplains ($n = 17$ spp., $r = 0.18$, $p = 0.071$), whereas more productive woodland ($n = 26$ spp., $p = 0.18$) and rainforest ($n = 15$ spp., $p = 0.94$) environments returned no significant relationship. The correlation between body mass dissimilarity and range overlap became statistically non-significant when controlling for phylogenetic distance (partial Mantel test, $p = 0.13$). Range overlap and dissimilarity in molar row length was also not significantly related (Mantel test, $p = 0.62$).

3.5 Local scale species co-occurrence analyses

Forty-one of the 55 named extant Australian dasyurid species were recorded at least once in the 83 local survey plots included in this study, with the antechinus *Antechinus stuartii*, and dunnarts *Sminthopsis crassicaudata* and *Sminthopsis dolichura* being represented in 24%, 23%, and 18% of the total sample set, respectively. Overall species richness (α -diversity) was low at most sites (average = 3.1 spp.), but this may reflect sampling biases resulting from more frequent documentation of mesic temperate regions in Western Australia, New South Wales, and Victoria. Conversely, the arid areas of South and Central Australia, and the tropical belt of northern Australia were under-represented, and it likely explains the conspicuous absence of the

desert-adapted mulgara, *Dasyercus blythi*, and tropical savannah inhabitants such as *Planigale ingrami*, from our dataset (Fig. 3). Absence of the Kakadu dunnart (*Sminthopsis bindi*), and near threatened Carpentarian pseudantechinus (*Pseudantechinus mimulus*) probably evidence their extremely restricted geographic distributions.

Of the 100 species spairs assessed using the methodology of Veech (2013), 22 showed significant levels of attraction, while only seven exhibited aversion (Table S4). Only one of the seven aversive (i.e., ‘checkerboard’) species pairs involved species with overlapping ranges: the antechinus *Antechinus flavipes*, and dunnart *S. crassicaudata*). All other pairs were allopatric species, specifically, these involved *A. stuartii* (endemic to eastern Australia, from southern Queensland to southern New South Wales) and the otherwise broadly distributed species of *Sminthopsis* and *Ningaui* (Fig. S2). Species pairs that showed a significant attraction were also not more morphologically divergent in body mass or molar row length compared with those expressing random or significant aversion (body mass: $t = 0.21$, $p = 0.83$; molar row length: $t = -0.23$, $p = 0.82$).

Lastly, we detected no meaningful association between the co-occurrence metric and extent of range overlap (Mantel test, $p = 0.40$: Fig. S3), or morphological dissimilarity (Mantel test, $p = 0.93$) when all geographically overlapping species pairs were examined. This result was consistent when body mass and molar row length were analyzed separately (all p -values >0.05). Nevertheless, the levels of co-occurrence did decrease with phylogenetic distance (Mantel test, $r = 0.07$, $p = 0.01$).

3.6 Node-based analysis

Only one node (marked “A” in Fig. 1) within our total evidence node + tip dated tree exhibited a high degree of allopatry (GND score >0.65), indicating distributional segregation amongst its descendent lineages. This corresponds to the late Miocene split (about 7.8 Ma) between the Australian wambengers (*Phascogale*) and New Guinean murexias (*Murexia*). Most other nodes with high GND scores were dispersed throughout the basal divergences in our topology, and constitute major clade subdivisions within Dasyuridae.

4. DISCUSSION

4.1 Sister species comparisons: testing for character displacement

The level of sympatry (52%) observed in our sample of dasyurid sister species pairs was slightly higher than that generally reported amongst placental mammals (33-45%: Fitzpatrick & Turelli, 2006). Our analysis of range overlap and trait divergence showed no evidence that competition reduces geographic range overlap in species with similar body sizes, or conversely, that range overlap promotes body size divergence. This lack of competitive displacement might be byproduct of climatic niche conservatism, which clearly affects climatic tolerances within Sminthopsini -one of the most speciose arid-adapted dasyurid clades- (García-Navas & Westerman, 2018). However, when testing if sister species were more similar to each other in terms of body mass and dentition with respect to other non-sister congeners, we found that it was true for roughly one quarter of cases (25% for body mass; 31% for molar row length). At this point, two important factors must be considered when interpreting these results. First, the existence of geographic range overlap does not guarantee co-occurrences at local scale. As our data shows (Table S3), species can distribute micro-allopatrically within local scale communities despite manifesting regional scale sympatry. This minimizes the need for competition-driven trait divergence, and could thus explain the lack of a relationship between regional range overlap and morphological divergence, as well as between range overlap and *c*-scores. Second, our estimates may average over remarkable intraspecific variability; despite a lack of difference in species-wide mean values, populations in close proximity may exhibit character displacement (Anacker & Strauss, 2014).

4.2 The geography of speciation

When plotting range overlap against node age for our 21 identified sister species pairs, we found that allopatry potentially played a more impactful role than sympatry in dasyurid speciation processes. This concurs with the established observation that smaller-bodied mammals

(including gophers, kangaroo rats, pocket mice and shrews among others) tend to undergo allopatric speciation (Fitzpatrick & Turelli, 2006); however, our results were not significantly supported (even following exclusion of *Murexia naso* + *Murexia melanurus*: slope = 0.11, intercept = -3.68, $F = 3.47$, $p = 0.07$; Fig. S4), probably because of small sample size. Here, we restricted our analyses to recent speciation events with differing relative divergence times (i.e., sister species comparisons) in order to mitigate some concerns regarding the lability and dynamic nature of species distributions (see e.g. Losos & Glor, 2003; Skeels & Cardillo, 2019). Such phenomena can manifest as trait divergences expressed across overlapping regions in the past, but subsequently exhibit minimal range overlap because of shifting ranges over time (evidenced *via* equivocal speciation history models: Fitzpatrick & Turelli, 2006). Conversely, recent simulations suggest that speciation history can produce detectable distributional and phylogenetic signals, even when species ranges have evolved substantially (Skeels & Cardillo, 2019). Accordingly, our results show that even in families in which species are highly mobile (see e.g., McLean et al., 2019) and there is a moderate proportion (>50%) of sympatric sisters, it is possible to find a subtle signature of allopatric speciation (positive ARC) supporting the view that sympatric speciation is usually rare among vertebrates (Phillimore et al., 2008).

4.3 Range overlap and morphological divergence

We failed to find a significant relationship between range overlap and morphological distance in dasyurid species, which indicates that biotic interactions have not played an important role in determining the spatial distribution of dasyurid species. Also, we did not find a relationship between the level of range overlap and dissimilarity in molar row length. This is in contrast with that reported by Davies et al. (2007) in a study across carnivores. Davies and colleagues found that species that differ more in carnassial tooth length overlap more in their geographical range. They also concluded that interspecific competition for food is a critical factor determining species co-existence in carnivores as upper carnassial length was better predictor of geographical range overlap than either canines or body size (Davies et al., 2007). However, they did not conduct analyses within each family due to small sample size and, in fact, two out of the

seven families analysed -Herpestidae and Procyonidae- did not show a positive trend between range overlap and divergence in carnassial length, which suggests that the reported trend seems not to be widespread within mammal families. Dasyurids lack specialized carnassials since they are mostly insectivorous and their diet includes a low proportion of vertebrates which largely varies among species (Dickman, 2014). Consequently, dental morphology is probably not subject to strong selection in this family of microcarnivorous

With regard to body mass, we found a negative relationship between this trait and range overlap (Fig. S5), implying that convergent selection has occurred in sympatry, while divergent selection has occurred in allopatry (i.e., abiotic factors are dominant). However, when cross-referenced against habitat, we only recovered a close-to-significant relationship for those species living in grasslands and sandplains, suggesting that such hostile environments likely promote character convergence. Notably, this concurs with previous studies of desert mammal communities, which tend to yield high species diversity relative to body size (see Shenbrot *et al.*, 1999 for a review). Co-existence between species of different body sizes has been attributed to competitive exclusion (Brown & Nicoletto, 1991), size-related niche segregation affecting preferred food resources (Brown & Lieberman, 1973; Alhajerj & Stepan, 2018), and/or contrasting spatial distributions (vis-à-vis dispersed *versus* grouped food resources: e.g., Hutto, 1978). In contrast, Australian arid zone dasyurid communities (which support up to eight–nine species) exhibit an opposing trend -convergent body size- that may be influenced by two complementary factors. Firstly, strong selection for an optimal body size that accommodates for both thermoregulatory (e.g., body size reduction as an advantage for harsh climates) and dispersive requirements (e.g., body size increase facilitating migration to utilize widely distributed food resources) in harsh environments. Secondly, reduced probability of interspecific encounters because of low population densities (a product of unpredictable food supplies) and/or strong micro-habitat selection. Overall, our results thus support the hypothesis that competition exerts only minimal constraint on regional scale species coexistence within Dasyuridae (Dickman, 2003).

4.4 Local scale species co-occurrence analyses

At local scale, we found no evidence that competition has determined spatial distribution between species within dasyurid local communities. Indeed, within our sampled survey plots the number of species pairs showing attraction was significantly greater than the number evincing aversion. Our results are therefore inconsistent with what would be expected according to the competitive exclusion hypothesis (i.e., a prevalence of negative interactions; Connor & Simberloff, 1979), and supports the null hypothesis that dasyurid species co-occurrence is random at population level. Furthermore, we tested whether the likelihood that the distribution of one species has been directly affected by the presence of other species increased as the degree of range overlap increases. Interestingly, we discovered that species with substantial overlap tended to have lower *c*-scores (i.e., higher randomness), although this relationship was not statistically significant. It suggests that niche-based processes (competition) have a negligible impact on dasyurid distributions. Co-occurrence data also revealed that dasyurid assemblages do not exhibit close packing. Consequently, sister species pairs were usually uncommon in local communities with less than half of the sympatric sister species pairs being recorded in the same survey plot (Table S3). Lastly, we didn't find a relationship between morphological divergence and co-occurrence at a local scale, which reinforce findings obtained at the regional scale (see above); biotic interactions don't seem to play an important role in shaping the distribution ranges composition of dasyurid assemblages.

4.5 Node-based analysis

Lastly, our node-based analysis pinpointed only a single node exhibiting strong distributional change (geographic node divergence GND score >0.65). This corresponds to the marked distributional segregation between the Australian *Phascogale* and New Guinean *Murexia*. The otherwise universally low GND scores calculated for other clades within Dasyuridae (indicating a dearth of geographically differentiated lineages in Australia *versus* New Guinea) concurs with the recognised biogeographical ubiquity of the group, and the hypothesis that colonization of

novel New Guinean habitats probably occurred opportunistically via multiple waves of migration commencing in the later Neogene (García-Navas et al., 2018).

4.6 Conclusions

Overall our results highlight the analytical advantages of examining species overlap at different spatial resolutions (Cardillo & Warren, 2016). Moreover, by integrating approaches used to infer broad-scale, evolutionary processes (phylogenetic comparative methods) and those commonly used to study ecological interactions at fine scales (community ecology), we show that it is possible to obtain a comprehensive understanding of factors driving species distributions. Specifically, we posit that geographic isolation arising from niche conservatism (as opposed to biotic interactions including competitive exclusion) has played a pivotal role in shaping the speciation patterns, and led to one of the most widespread endemic mammal radiations found on the Australasian landmasses today (García-Navas & Westerman, 2018).

Acknowledgements

V. García-Navas is supported by a Forschungskredit from the University of Zurich (grant ref. FK-18-104). Andrew Baker (Queensland University of Technology, Australia) provided measurement data. Fabricio Villalobos (INECOL, Mexico) and two anonymous referees provided constructive comments that improved an earlier version of the manuscript.

REFERENCES

- Alhajeri, B.H. & Steppan, S.J. (2018) Community structure in ecological assemblages of desert rodents. *Biological Journal of the Linnean Society*, 124, 308-318.
- Anacker, B.L. & Strauss, S.Y. (2014) The geography and ecology of plant speciation: range overlap and niche divergence in sister species. *Proceedings of the Royal Society of London B*, 281, 20132980.
- Anderson, B.M., Thiele, K.R., Grierson, P.F., Krauss, S.L., Nevill, P.G., Small, I.D., Zhong, X. & Barrett, M.D. (2019) Recent range expansion in Australian hummock grasses (*Triodia*) inferred using genotyping-by-sequencing. *AoB Plants*, *in press*.

543 Archer, M. (1976) The dasyurid dentition and its relationships to that of didelphids, thylacinids,
 544 borhyaenids (Marsupicarnivora) and peramelids (Peramelina: Marsupialia). Australian
 545 Journal of Zoology, Supplementary Series, 39, 1-34.

546 Archer, M. (1978) The nature of the molar-premolar boundary in marsupials and a
 547 reinterpretation of the homology of marsupial cheekteeth. Memoirs of the Queensland
 548 Museum, 18, 157-164.

549 Borregaard, M.K., Rahbek, C., Fjeldsa, J., Parra, J.L., Whittaker, R.J. & Graham, C.H. (2014)
 550 Node-based analysis of species distributions. Methods in Ecology and Evolution, 5, 1225-
 551 1235.

552 Brown, J.H. & Lieberman, G.A. (1973). Resource utilization and coexistence of seed-eating
 553 desert rodents in sand dune habitats. Ecology, 54: 788-797.

554 Brown, J.H. & Nicoletto, P.F. (1991) Spatial scaling of species composition: body masses of
 555 North American land mammals. American Naturalist, 138: 1478-1512.

556 Cantalapiedra, J.L., Fernández, M.H. & Morales, J. (2014). The biogeographic history of
 557 ruminant faunas determines the phylogenetic structure of their assemblages at different
 558 scales. Ecography, 37, 1-9.

559 Cardillo, M. (2011) Phylogenetic structure of mammal assemblages at large geographical
 560 scales: linking phylogenetic community ecology with macroecology. Philosophical
 561 Transactions of the Royal Society of London B, 366, 2545-2553

562 Cardillo, M. (2012) The phylogenetic signal of species co-occurrence in high-diversity
 563 shrublands: different patterns for fire-killed and fire-resistant species. BMC Ecology, 12, 1-
 564 10.

565 Cardillo, M. & Warren, D.L. (2016) Analysing patterns of spatial and niche overlap among
 566 species at multiple resolutions. Global Ecology and Biogeography, 25, 951-963.

567 Catullo, R.A. & Keogh, J.S. (2014) Aridification drove repeated episodes of diversification
 568 between Australian biomes: evidence from a multi-locus phylogeny of Australian toadlets
 569 (Uperoleia: Myobatrachidae). Molecular Phylogenetics and Evolution, 79, 106-117.

570 Cavender-Bares, J., Kozak, K.H., Fine, P.V.A. & Kembel, S.W. (2009) The merging of
 571 community ecology and phylogenetic biology. Ecology Letters, 12, 693-715.

572 Chesser, R. T. & Zink, R.M. (1994) Modes of speciation in birds - a test of Lynch's method.
 573 Evolution, 48, 490-497.

574 Cody, M.L. (1969) Convergent characteristics in sympatric species: a possible relation to
 575 interspecific competition and aggression. Condor, 71, 223-239.

576 Connor EF, Simberloff D (1979) The assembly of species communities: chance or competition?
 577 Ecology, 60, 1132-1140.

578 Crowther, M. & Blacket, M. (2003) Biogeography and speciation in the Dasyuridae: why are
 579 there so many kinds of Dasyurids? In Jones, M.E., Dickman, C.R. & Archer, M. (Eds.),

580 Predators with Pouches, (pp. 124-130). Australia: CSIRO Publishing.
 581 Davies, T.J., Meiri, S., Barraclough, T.G. & Gittleman, J.L. (2007) Species co-existence and
 582 character divergence across carnivores. *Ecology Letters*, 10, 146-152.
 583 Dickman, C.R. (1984) Competition and coexistence among the small marsupials of Australia
 584 and New Guinea. *Acta Zoologica Fennica*, 172, 27-31.
 585 Dickman, C.R. (2003) Distributional ecology of Dasyurid marsupials. In: Jones, M.E.,
 586 Dickman, C.R. & Archer, M. (Eds.), *Predators with Pouches*, (pp. 318-331). Australia:
 587 CSIRO Publishing.
 588 Dickman, C.R. (1986a) An experimental study of competition between two species of dasyurid
 589 marsupials. *Ecological Monographs*, 56, 221-241.
 590 Dickman, C.R. (1986b) Niche compression: two tests of an hypothesis using narrowly
 591 sympatric predator species. *Australian Journal of Ecology*, 11, 121-134.
 592 Dickman, C.R. (1988) Body size, prey size, and community structure in insectivorous
 593 mammals. *Ecology*, 69, 569-580.
 594 Dickman, C.R. (2014) Micro-carnivores: the ecological role of small-dasyurid predators in
 595 Australia. In: Glen, A. & Dickman, C.R. (eds.) *Carnivores of Australia: past, present, and*
 596 *future*, (pp. 241-262). Australia: CSIRO Publishing.
 597 Drury, J.P., Tobias, J.A., Burns, K.J., Mason, N.A., Shultz, A.J. & Morlon, H. (2018)
 598 Contrasting impacts of competition on ecological and social trait evolution in songbirds.
 599 *PLoS Biology*, 16, e2003563.
 600 Felsenstein, J. (1985) Phylogenies and the comparative method. *American Naturalist*, 125, 1-15.
 601 Fisher, D.O. & Dickman, C.R. (1993) The body size-prey relationship in dasyurid marsupials:
 602 tests of three hypotheses. *Ecology*, 74, 1871-1883.
 603 Fisher, D.O. & Cockburn, A. (2006) The large-male advantage in brown antechinuses:
 604 female choice, male dominance, and delayed male death. *Behavioral Ecology*, 17, 164-171.
 605 Fitzpatrick, B.M. & Turelli, M. (2006) The geography of mammalian speciation: mixed signals
 606 from phylogenies and range maps. *Evolution*, 60, 601-615.
 607 Fox, B.J. (1982) A review of dasyurid ecology and speculation on the role of limiting similarity
 608 in community organization. In: Archer, M. (Eds.), *Carnivorous marsupials*, (pp. 318-331).
 609 Australia: CSIRO Publishing.
 610 Gaboriau, T., Leprieux, F., Mouillot, D. & Hubert, N. (2018) Influence of the geography of
 611 speciation on current patterns of coral reef fish biodiversity across the Indo-Pacific.
 612 *Ecography*, 41: 1295-1306.
 613 García-Navas, V. & Westerman, M. (2018) Niche conservatism and phylogenetic clustering in a
 614 tribe of arid-adapted marsupial mice, the Sminthopsini. *Journal of Evolutionary Biology*, 31:
 615 1204-1215.

616 García-Navas, V., Rodríguez-Rey, M. & Westerman, M. (2018) Bursts of morphological and
617 lineage diversification in modern dasyurids, a “classic” adaptive radiation. *Biological Journal*
618 of the Linnean Society, 123, 892-795.

619 Gordon, C.L. (2003) A first look at estimating body size in dentally conservative marsupials.
620 *Journal of Mammalian Evolution*, 10, 1-21.

621 Grant, P. & Grant, R. (2006) Evolution of character displacement in Darwin’s finches. *Science*,
622 313, 224-226.

623 Griffith, D.M., Veech, J.A. & Marsh, C.J. (2016) *cooccur*: Probabilistic Species Co-Occurrence
624 Analysis in R. *Bioinformatics*, 69, code snippet 2.

625 Grossenbacher, D.L. & Whittall, J.B. (2011) Increased floral divergence in sympatric
626 monkeyflowers. *Evolution*, 65, 2712-2718.

627 Hodgell, J.R. & Bellwood, D.R. (2016) The geography of speciation in coral reef fishes: the
628 relative importance of biogeographical barriers in separating sister-species. *Journal of*
629 Biogeography, 43, 1324-1335.

630 Hutto, R.L. (1978) A mechanism for resource allocation among sympatric rodent species.
631 *Oecologia*, 33: 115-126.

632 IUCN (2018) The IUCN Red List of Threatened Species. Version 2018-2. <[http://](http://maps.iucnredlist.org)
633 maps.iucnredlist.org>. Accessed on: September 2018.

634 Jombart, T., Balloux, F. & Dray, S. (2010) *adephylo*: new tools for investigating the
635 phylogenetic signal in biological traits, *Bioinformatics*, 26, 1907-1909.

636 Jones, M.E. (1997) Character displacement in Australian dasyurid carnivores: size relationships
637 and prey size patterns. *Ecology*, 78, 2569-2587.

638 Jones, M.E. & Barmuta, L.A. (2000) Niche differentiation among sympatric Australian dasyurid
639 carnivores. *Journal of Mammalogy* 81, 434-447.

640 Losos, J.B. & Glor, R.E. (2003) Phylogenetic comparative methods and the geography of
641 speciation. *Trends in Ecology and Evolution*, 18, 220-227.

642 Kealy, S. & Beck, R (2017) Total evidence phylogeny and evolutionary timescale for Australian
643 faunivorous marsupials (Dasyuromorphia). *BMC Evolutionary Biology*, 17, 240.

644 Kear, B.P., Aplin, K.P. & Westerman, M. (2016) Bandicoot fossils and DNA elucidate lineage
645 antiquity amongst xeric-adapted Australasian marsupials. *Scientific Reports*, 6, 37537.

646 Kembel, S.W., Cowan, P.D., Helmus, M.R., Cornwell, W.K., Morlon, H., Ackerly, D.D.,
647 Blomberg, S.P. & Webb, C.O. (2010). *picante*: R Tools for Integrating Phylogenies and
648 Ecology. *Bioinformatics*, 26, 1463.

649 Krajewski, C., Wroe, S. & Westerman, M. (2000) Molecular evidence for the pattern and timing
650 of cladogenesis in dasyurid marsupials. *Zoological Journal of the Linnean Society*, 130, 375-
651 404.

652 Kraaijeveld-Smit, F.J.L., Ward, S.J. & Temple-Smith, P.D. (2003) Paternity success and the
653 direction of sexual selection in a field population of a semelparous marsupial, *Antechinus*
654 *agilis*. *Molecular Ecology*, 12, 475-484.

655 MacArthur, R. & Levins, R. (1967) The limiting similarity, convergence, and divergence of
656 coexisting species. *The American Naturalist*, 101, 377-385.

657 Martin, S.A., Alhajeri, B.H. & Steppan, S.J. (2016) Dietary adaptations in the teeth of murine
658 rodents (Muridae): a test of biomechanical predictions. *Biological Journal of the Linnean*
659 *Society*, 119, 766-784.

660 Martin, P.R., Montgomerie, R. & Loughheed, S.C. (2010) Rapid sympatry explains greater color
661 pattern divergence in high latitude birds. *Evolution*, 64, 336-347.

662 McLean, A.L., Cooper, S.J.B., Lancaster, M.L., Gaikhorst, G., Lambert, C., Moseby, K., Read,
663 J., Ward, M. & Carthew, S.M. (2019) Small marsupial, big dispersal? Broad- and fine-scale
664 genetic structure of an endangered marsupial from the Australian arid zone. *Australian*
665 *Journal of Zoology*, 66, *in press*.

666 Morton, S.R., Dickman, C.R. & Fletcher, T.P. (1989) *Dasyuridae*. In: *Fauna of Australia vol. 1B*
667 *Mammalia*. Walton, D.W. & Richardson, B.J. (eds.) AGPS, Canberra.

668 Mouquet, N., Devictor, V., Meynard, C.N., Munoz, F., Bersier, L.-F., Chave, J., Couteron, P.,
669 Dalecky, A., Fontaine, C., Gravel, D., Hardy, O.J., Jabot, F., Lavergne, S., Leibold, M.,
670 Mouillot, D., Münkemüller, T., Pavoine, S., Prinzing, A., Rodrigues, A.S.L., Rohr, R.P.,
671 Thébault, E. & Thuiller, W. (2012) Ecophylogenetics: advances and perspectives. *Biological*
672 *Reviews*, 87, 769-785.

673 Oksanen, J., Kindt, R., Legendre, P., & O'Hara, R.B. (2010) *vegan*: community ecology
674 package. Version 1.7–81 (<http://cran.r-project.org/>).

675 Paradis, E., Claude, J. & Strimmer, K. (2004) *APE*: Analyses of Phylogenetics and Evolution in
676 R language, *Bioinformatics*, 20: 289-290.

677 Phillimore, A. B., Orme, C.D.L., Thomas, G.H., Blackburn, T.M., Bennett, P.M., Gaston, K.J.
678 & Owens, I.P.F. (2008) Sympatric speciation in birds is rare: insights from range data and
679 simulations *American Naturalist*, 171: 646-657.

680 Pineda-Munoz, S., Lazagabaster, I.A., Alroy, J. & Evans, A.R. (2017), Inferring diet from
681 dental morphology in terrestrial mammals. *Methods in Ecology and Evolution*, 8, 481-491.

682 Rabosky, D.L., Donnellan, S.C., Talaba, A.L. & Lovette, I.J. (2007) Exceptional among-lineage
683 variation in diversification rates during the radiation of Australia's most diverse vertebrate
684 clade. *Proceedings of the Royal Society of London B*, 274, 2915-2923.

685 Shenbrot, G.I., Krasnov, B. & Rogovin, K.A. (1999) *Spatial Ecology of Desert Rodent*
686 *Communities*. Springer Verlag, Berlin.

687 Stone, L. & Roberts, A. (1990) The checkerboard score and species distributions. *Oecologia*,

85: 74-79.

- Travouillon, K. & Philips, M.J. (2018) Total evidence analysis of the phylogenetic relationships of bandicoots and bilbies (Marsupialia: Paramelomorphia): Reassessment of two species and description of a new species. *Zootaxa*, 4378, 224.
- Vamosi, S.M., Heard, S.B., Vamosi, J.C. & Webb, C.O. (2009) Emerging patterns in the comparative analysis of phylogenetic community structure. *Molecular Ecology*, 18, 572-592.
- van Dyck, S. & Strahan, S. (2008) *The Mammals of Australia*. New Holland publishers, Sydney.
- Veech, J.A. (2013) A probabilistic model for analysing species co-occurrence: probabilistic model. *Global Ecology and Biogeography*, 22, 252–260.
- Vilela, B. & Villalobos, F. (2015) *letsR*: a new R package for data handling and analysis in macroecology. *Methods in Ecology and Evolution*, 6: 1229-1234.
- Warren, D.L., Cardillo, M., Rosauer, D.F. & Bolnick, D.I. (2014) Mistaking geography for biology: inferring processes from species distributions. *Trends in Ecology and Evolution*, 29, 572-580.
- Weeks, B.C., Claramunt, S. & Cracraft, J. (2016) Integrating systematics and biogeography to disentangle the roles of history and ecology in biotic assembly. *Journal of Biogeography*, 43, 1546-1559.
- Webb, C.O., Ackerly, D.D., McPeck, M.A. & Donoghue, M.J. (2002) Phylogenies and community ecology. *Annual Review of Ecology and Systematics*, 33, 475-505.
- Westerman, M., Krajewski, C., Kear, B.P., Meehan, L., Meredith, R.W., Emerling, C.A. & Springer, M.S. (2016) Phylogenetic relationships of dasyuromorphian marsupials revisited. *Zoological Journal of the Linnean Society*, 176, 686-701.
- Westerman, M., Blacket, M.J., Hintz, A., Armstrong, K., Woolley, P.A. & Krajewski, C. (2017) A plethora of planigales: genetic variability and cryptic species in a genus of dasyurid marsupials from northern Australia. *Australian Journal of Zoology*, 64, 303-311.
- Wiens, J. (2012) Why biogeography matters: historical bio-geography vs. phylogeography and community phylogenetics for inferring ecological and evolutionary processes. *Frontiers of Biogeography*, 4, 128-135.
- Woolley, P.A. (1992) New records of the Julia Creek dunnart, *Sminthopsis douglasi* (Marsupialia: Dasyuridae). *Wildlife Research*, 19, 779-783.

Biosketch

Vicente García-Navas is a postdoctoral fellow at the University of Zurich in Switzerland. His research focuses on the macroevolutionary implications of geographic diversity concentrations and niche conservatism in vertebrates.

Benjamin P. Kear is Docent and Curator of Vertebrate Palaeontology at the Museum of Evolution in Uppsala University, Sweden, and researches the deep time evolution of australidelphian marsupials.

Michael Westerman is an Adjunct Professor at La Trobe University in Australia, and a leading specialist on the genetics and molecular systematics of Australasian mammals.

Author contributions

V.G.N. designed the study; V.G.N and M.W. compiled the ecomorphological data; M.W. and B.P.K. compiled the phylogenetic data and analyses; V.G.N carried out the analyses and wrote the manuscript with input from B.P.K. and M.W. All authors read and approved the final manuscript.

Data availability statement

The data supporting the results are available from the LabArchives Repository:
XXXXXXXXXXXXXXXXXX

Title: Data from: The geography of speciation in dasyurid marsupials

DOI: XXXXXXXXXXXXXXXX

Journal: Journal of Biogeography

Journal manuscript number: JBI-19-XXXX

Fig. 1. Time-calibrated maximum clade credibility tree of crown Dasyuridae. Blue and red branches represent sympatric and allopatric sister species, respectively. Sister species pairs with posterior probability < 1 are indicated with an asterisk. Gradient scale (top right) and node symbol diameter (coloured circles) are proportional to GND at each node (see main text). Only fully resolved nodes where both descendant clades consist of at least two species were included in the node-based analysis. Distributional maps (“A-C”) represent SOS values for the corresponding nodes (“A-C”) with high GND scores; highest GND score > 0.65 (arrow). Colour gradient indicates over-representation of opposing descendant clades (red versus blue) *versus* both descendant clades being equally represented (yellow).

Fig. 2. Scatter plot of (transformed) range overlap against body mass divergence quantified as morphological distance (square-root transformed Euclidean distance) based on pairwise differences in body mass (Mantel test, $r = 0.06$, $p = 0.015$). In the x-axis, a value of zero means complete overlap (i.e., complete sympatry), and a value of 1 would mean complete allopatry. Both sympatric and allopatric pairs of species are included. Exclusion of non-overlapping pairs resulted in a steeper linear relationship (grey regression line; see Supplementary Material). Artwork courtesy Kevin Stead.

Fig. 3. Map of Australia showing the approximate location of the 83 local survey plots (representing small mammal communities) herein (see Supplementary Material). Pie charts indicate the percentage of n -species communities in each region. Richer communities (> 6 species) are located in the arid areas of Queensland, northeastern Australia. The map of the inset shows the spatial distribution of species richness of dasyurid marsupials in the Australo-Papuan region. The Australian arid zone is the region with the greatest species diversity (yellow-orange areas) whereas some regions across the ‘Top End’, Northern Territory (empty cells), do not harbor any dasyurid species.

772

773

774

775





Figure 2



790

